

5223.0080 MUSCULOSKELETAL SCHEDULE; AMPUTATIONS OF UPPER EXTREMITY.

Permanent partial disability due to amputation of upper extremities is a disability of the whole body as follows:

- A. forequarter amputation, 70 percent;
- B. disarticulation at shoulder joint, 60 percent;
- C. amputation of arm above deltoid insertion, 60 percent;
- D. amputation of arm between deltoid insertion and elbow joint, 57 percent;
- E. disarticulation at elbow joint, 57 percent;
- F. amputation of forearm below elbow joint proximal to insertion of biceps tendon, 57 percent;
- G. amputation of forearm below elbow joint distal to insertion of biceps tendon, 54 percent;
- H. disarticulation at wrist joint, 54 percent;
- I. midcarpal or midmetacarpal amputation of hand, 54 percent;
- J. amputation of all fingers except thumb at metacarpophalangeal joints, 32.5 percent;
- K. amputation of thumb:
 - (1) at metacarpophalangeal joint or with resection of metacarpal bone, 21.5 percent;
 - (2) at interphalangeal joint or through proximal phalanx, 16 percent;
 - (3) from interphalangeal joint to midportion distal phalanx, 13 percent;
 - (4) from middistal phalanx, distal, 6 percent;
- L. amputation of index finger:
 - (1) at metacarpophalangeal joint or with resection of metacarpal bone or through proximal phalanx, 13.5 percent;
 - (2) at proximal interphalangeal joint or through middle phalanx, 11 percent;
 - (3) at distal interphalangeal joint to middistal phalanx, 5 percent;
 - (4) from middistal phalanx, distal, 2.5 percent;

M. amputation of middle finger:

- (1) at metacarpophalangeal joint or with resection of metacarpal bone or through proximal phalanx, 11 percent;
- (2) at proximal interphalangeal joint or through middle phalanx, 9 percent;
- (3) at distal interphalangeal joint to middistal phalanx, 5 percent;
- (4) from middistal phalanx, distal, 2.5 percent;

N. amputation of ring finger:

- (1) at metacarpophalangeal joint or with resection of metacarpal bone or through proximal phalanx, 5.5 percent;
- (2) at proximal interphalangeal joint or through middle phalanx, 4 percent;
- (3) at distal interphalangeal joint to middistal phalanx, 3 percent;
- (4) from middistal phalanx, distal, 1.5 percent;

O. amputation of little finger:

- (1) at metacarpophalangeal joint or with resection of metacarpal bone or through proximal phalanx, 3 percent;
- (2) at proximal interphalangeal joint or through middle phalanx, 2 percent;
- (3) at distal interphalangeal joint to middistal phalanx, 1 percent;
- (4) from middistal phalanx, distal, 0.5 percent.

Statutory Authority: *MS s 176.105*

History: *10 SR 1124*

Published Electronically: *August 16, 2010*